



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

PROBLEMS FOR SOLUTION.

ALGEBRA.

170. Proposed by S. F. NORRIS, Professor of Astronomy and Mathematics, Baltimore City College, Baltimore, Md.

Find by strictly quadratic methods at least one set of values of x and y in the equations $x^2y^2 + x = 38$ and $xy + y^2 = 15$.

171. Proposed by IDA M. SCHOTTENFELTZ, A. M.

$ay^2 + a = bxy + cx$, $bx^2 + b = axy + cy$. Solve for x and y .

172. Proposed by L. E. DICKSON, Ph. D., The University of Chicago.

Without solving the algebraically solvable quintic $y^5 + py^3 + \frac{1}{5}p^2y + r = 0$, prove that it is irreducible in the domain of rationality (p, r) .

GEOMETRY.

193. Proposed by PROFESSOR BEYENS.

Si le rapport du segment d'une base de la sphère à l'hémisphère est m/n , le rapport de l'hauteur du segment à deux bases qui resultera au rayon est égal à $2\sin\frac{1}{2}[\sin^{-1}(n-m)/n]$. [Problem 9699, *Educational Times*.]

194. Proposed by MARCUS BAKER, U. S. Geological Survey, Washington, D. C.

Glass paper weights, having the form of a regular tetrahedron, are to be packed for shipment, each in a paper box. Wanted to know the size and shape of the smallest box for the purpose. How much empty space in each box?

CALCULUS.

160. Proposed by B. F. FINKEL, A.M., M.Sc., Professor of Mathematics and Physics, Drury College, Springfield, Mo.

A dog at the vertex of a right conical hill pursues a fox at the foot of the hill. How far will the dog run to catch the fox, if the dog runs directly toward the fox at all times, and the fox is continually running around the hill at its foot, the velocity of the dog being 6 feet per second, the velocity of the fox being 5 feet per second, the hill being 100 feet high and 200 feet in diameter at the base?

MECHANICS.

150. Proposed by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

O is a point in the plane of a triangle, ABC , and D, E, F are the mid-points of the sides. Show, geometrically, that the system of forces OA, OB, OC is equivalent to the system OD, OE, OF .